

Normansfield Statistics 1868—1886

<u>Year</u>	<u>Number Admitted</u>	<u>Deaths</u>	<u>Discharges</u>	<u>Total Number</u>	<u>Average Age on Admission</u>
1868	21	none	2	19	13.4
1869	5	none	1	22	9.0
1870	13	1	2	30	13.0
1871	13	none	7	37	15.5
1872	13	none	2	49	13.0
1873	11	2	3	57	11.4
1874	27	8	none	75	14.9
1875	24	3	8	90	12.5
1876	17	4	15	93	11.8
1877	25	5	8	100	11.6
1878	19	3	8	106	13.0
1879	35	8	10	117	10.2
1880	15	2	10	117	12.3
1881	30	3	10	129	12.6
1882	25	1	11	145	10.1
1883	13	6	14	145	13.0
1884	24	2	12	149	11.6
1885	7	2	15	143	10.0
1886	13	5	18	139	13.0

Causes of Death in Normansfield 1868—1886

The death rates could be ascribed in the main to infection and to epilepsy. The distribution of causes of death is set out below. The figures are drawn from the review of medical records.

Total Deaths	55
Mean annual death rate	2.8%
Tuberculosis	3
Epilepsy	12
Scarlet Fever	8
Measles	4
Respiratory infection	14
Others	14

Statistical extracts from PhD Thesis: Prof Conor Ward 1999